


Accepted Manuscript

Duration of Dual Antiplatelet Therapy Following Drug-Eluting Stent Implantation in Diabetic and Non-Diabetic Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials

Abhishek Sharma, Aakash Garg, Sammy Elmariah, Douglas Drachman, Chukwudi Obiagwu, Ajay Vallakati, Samin K. Sharma, Carl J. Lavie, Debabrata Mukherjee, Ron Waksman, Giulio G. Stefanini, Fausto Feres, Jonathan D. Marmur, Gérard Helft



PII: S0033-0620(17)30165-2
DOI: <https://doi.org/10.1016/j.pcad.2017.12.003>
Reference: YPCAD 854

To appear in:

Received date: 17 December 2017
Accepted date: 17 December 2017

Please cite this article as: Abhishek Sharma, Aakash Garg, Sammy Elmariah, Douglas Drachman, Chukwudi Obiagwu, Ajay Vallakati, Samin K. Sharma, Carl J. Lavie, Debabrata Mukherjee, Ron Waksman, Giulio G. Stefanini, Fausto Feres, Jonathan D. Marmur, Gérard Helft, Duration of Dual Antiplatelet Therapy Following Drug-Eluting Stent Implantation in Diabetic and Non-Diabetic Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Ypcad(2017), <https://doi.org/10.1016/j.pcad.2017.12.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Duration of Dual Antiplatelet Therapy Following Drug-Eluting Stent Implantation in Diabetic and Non-Diabetic Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials

Abhishek Sharma^{1,2}, MD; Aakash Garg², MD Sammy Elmariah^{1,4}, MD, MPH; Douglas Drachman¹, MD; Chukwudi Obiagwu⁵, MD;; Ajay Vallakati⁶, MD; Samin K Sharma⁷, MD; Carl J Lavie⁸, MD; Debabrata Mukherjee⁹, MD, MS; Ron Waksman¹⁰, MD; Giulio G. Stefanini¹¹, MD, Ph.D; Fausto Feres¹², MD, PhD; Jonathan D. Marmur¹³, MD; Gérard Helft^{14,15}, MD

1. Division of Cardiovascular Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA
2. Institute of Cardiovascular Science and Technology, Brooklyn, NY, USA
3. Division of Cardiology, Newark Beth Israel Medical Center, Newark, NJ, USA
4. Harvard Clinical Research Institute, Boston, USA
5. Department of Cardiovascular Medicine, Maimonides Medical Center, Brooklyn, NY, USA
6. Division of Cardiology, Metrohealth Medical Center, Case Western Reserve University, Cleveland, OH, USA
7. Department of Cardiovascular Medicine, Heart & Vascular Institute, Mount Sinai Medical Center, Icahn School of Medicine at Mount Sinai, New York, NY, USA
8. Department of Cardiovascular Diseases, John Ochsner Heart and Vascular Institute, Ochsner Clinical School-the University of Queensland School of Medicine, New Orleans, LA, USA
9. Division of Cardiology, Texas Tech University, El Paso, TX, USA
10. Division of Cardiology, MedStar Washington Hospital Center, Washington, DC, USA

11. Cardiovascular Department, Humanitas Research Hospital, Rozzano, Milan, Italy
12. Instituto Dante Pazzanese de Cardiologia, Ave Dante Pazzanense, 500,
Ibirapuera, São Paulo, São Paulo, Brazil
13. Division of Cardiovascular Medicine, State University of New York, Downstate
Medical Center, Brooklyn, NY, USA
14. Institut de Cardiologie, Hôpital Pitié-Salpêtrière, Assistance Publique Hôpitaux de
Paris, Université Pierre et Marie Curie, boulevard de l'Hôpital, Paris, France
15. Institute of Cardiometabolism and Nutrition, Hôpital Pitié-Salpêtrière, Paris,
France

Corresponding Author:

Abhishek Sharma, MD

Postal address: Division of Cardiovascular Medicine

Massachusetts General Hospital and Harvard Medical School

55 Fruit St, Boston, MA 02114

Phone# 201-892-6548

Fax: 718-270-2917

Email: abhisheksharma4mamc@gmail.com

Short title: DAPT after DES in Diabetic and Non-Diabetic Patients

Disclosure: None

ABSTRACT:

Background: Diabetic patients account for an increasing number of patients undergoing percutaneous coronary intervention (PCI). However, diabetes mellitus (DM) is associated with increased residual platelet activity during dual antiplatelet treatment (DAPT) and DM patients have worse clinical outcomes after PCI as compared to non-DM.

Objective: To evaluate efficacy and safety of short duration DAPT (S-DAPT) and long duration DAPT (L-DAPT) after drug eluting stent (DES) implantation in DM and non-DM patients.

Methods: We searched Medline, Embase, and Cochrane Central Register of Controlled Trials (CENTRAL) to identify randomized controlled trials (RCTs) assessing the effect of S-DAPT versus L-DAPT after DES implantation in DM and non-DM patients.

Efficacy endpoints were all-cause mortality, cardiac mortality, myocardial infarction (MI), stent thrombosis (ST), target vessel revascularization (TVR), and composite end point of net adverse clinical events (NACE) (all cause mortality, cardiac mortality, MI, ST, TVR, stroke, major bleeding). Safety endpoints were major bleeding and stroke. Event rates were compared using a forest plot of relative risk using a random effects model.

Results: We included eight RCTs that randomized 28,318 patients to S-DAPT versus L-DAPT (8,234 DM and 20,084 non-DM). S-DAPT was associated with an increased rate of ST in non-DM patients [3.67 (2.04, 6.59)]. There was no significant difference in the rate of all cause mortality, cardiac mortality, ST, MI, TVR, major bleeding, stroke and NACE with S-DAPT and L-DAPT in DM patients [1.19 (0.72-1.95); 1.25 (0.69, 2.25);

1.52 (0.70, 3.29); 1.33 (0.88, 2.01); 1.39 (0.89, 2.17); 0.92 (0.19, 4.42); 0.98 (0.29, 3.28); and 0.94 (0.57, 1.54) respectively]. Further, there was no significant difference in the rate of all cause mortality, cardiac mortality, MI, TVR, major bleeding, stroke and NACE with S-DAPT and L-DAPT in non-DM patients [0.93 (0.58, 1.48); 0.75 (0.42, 1.35); 1.52 (0.81, 2.83); 0.99 (0.71, 1.39); 0.72 (0.28, 1.84); 1.01 (0.40, 2.56); and 1.01 (0.77, 1.32) respectively].

Conclusion: Compared to L-DAPT, S-DAPT was associated with significant increase in rate of ST in non-DM patients. Duration of DAPT had no significant impact on rates of all cause mortality, cardiac mortality, MI, ST and TVR among DM patients.

Key Words: Dual anti-platelet therapy, Drug eluting stents, Diabetes

ABBREVIATIONS

CAD: Coronary Artery Disease

DAPT: Dual Anti-platelet Therapy

DES: Drug Eluting Stent

DM: Diabetes Mellitus

L-DAPT: Long Dual Anti-platelet Therapy

MI: Myocardial Infarction

NACE: Net Adverse Cardiovascular Event

PCI: Percutaneous Coronary Intervention

RCT: Randomized Controlled Trial

S-DAPT: Short Dual Anti-platelet Therapy

ST: Stent Thrombosis

TVR: Target Vessel Revascularization

INTRODUCTION:

The prevalence of diabetes mellitus (DM) is increasing globally, and optimal care of DM patients remains a challenge for health care providers. Patients with DM have higher prevalence of coronary artery disease (CAD) and account for an increasing number of patients undergoing percutaneous coronary intervention (PCI) [1]. However, DM patients have worse clinical outcomes after PCI as compared to non-DM [1, 2].

Dual antiplatelet therapy (DAPT) using a combination of aspirin and a P2Y₁₂ inhibitor is used for the prevention of ischemic complications after PCI. However, the optimal duration of DAPT after drug-eluting stent(s) (DES) implantation remains unclear [3]. Several studies have reported an increase in bleeding complications with longer duration (>12 months) of DAPT (L-DAPT) compared to short duration (\leq 12 months) DAPT (S-DAPT), without any significant improvement in the rates of atherothrombotic complications [4-10]. However, the large multicenter DAPT Study recently demonstrated improved clinical outcomes with longer duration of DAPT (L-DAPT) compared to short duration DAPT (S-DAPT) [11]. Pooled analyses of various randomized controlled trials (RCTs) have reported conflicting results [12-20]. Inconsistent results are likely due to the evaluation of different studies and patient subgroups. Diabetes mellitus is associated with a prothrombotic state secondary to endothelial dysfunction, hyper-coagulability, and platelet activation. Thus, the benefits of L-DAPT versus S-DAPT in diabetic patients warrant investigation. Against this background, we performed a meta-analysis of RCTs to

evaluate the efficacy and safety of S-DAPT and L-DAPT after DES implantation in DM and non-DM patients.

METHODS:

We followed the PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) statement for reporting systematic reviews and meta-analyses of RCTs for the protocol of our meta-analysis [21].

We systematically searched PubMed, CINAHL, Cochran CENTRAL, Embase, Scopus and Web of Science databases for randomized clinical trials comparing different durations of DAPT after DES implantation in patients with and without DM. Pertinent trials were also searched in clinicaltrials.gov and in the proceedings of major international cardiology meetings (ACC, AHA, ESC, The Society for Cardiac Angiography and Interventions). In addition, the reference lists of the identified studies were searched for any potential relevant studies. DAPT was defined as aspirin plus a P2Y₁₂ receptor inhibitor, after coronary DES implantation. S-DAPT and L-DAPT were defined as duration of DAPT after DES implantation ≤ 12 months and > 12 months respectively. All relevant combinations of following keywords “aspirin”, “P2Y₁₂ receptor inhibitor”, “clopidogrel”, “Plavix”, “prasugrel”, “Effient”, “ticagrelor”, “Brilinta”, “thienopyridine”, “dual antiplatelet therapy”, “DAPT”, “drug eluting stents”, “DES”, “diabetes mellitus”, “DM”, “non diabetics”, “non DM”, “death”, “mortality”, “survival”, “cardiac mortality”, “stent thrombosis”, “Global Utilization Of Streptokinase

And Tpa For Occluded Arteries (GUSTO) bleeding”, stroke, myocardial infarction, “randomized controlled trial”, “random”, “random allocation”, “double-blind”, and “single-blind” were included for database search. We manually searched references of identified studies. Only studies published between January 1, 2002 to June 30, 2016 were included. No language restrictions were applied. Studies, which did not report the absolute numbers of events in DM and non-DM patients after DES implantation, were excluded from the analysis. The corresponding authors of the relevant studies were queried for required quantitative details not in the published manuscripts.

Two reviewers (AS, AG) independently screened the titles and abstracts for relevance. The full texts of selected manuscripts were reviewed for inclusion or exclusion using the above mentioned selection criteria. Two reviewers (AS, AG) independently determined the articles to be included and excluded, and data from the relevant articles were extracted using pre-defined extraction forms. Any disagreements in data extraction were discussed until consensus was reached. Baseline patient characteristics and follow up data was extracted from selected studies at the end of follow up period: all-cause mortality, cardiac mortality, myocardial infarction (MI), stent thrombosis (ST), target vessel revascularization (TVR), major bleeding, stroke and net adverse cardiovascular event (NACE), the composite of all cause mortality, cardiac mortality, MI, ST, TVR, stroke, major bleeding. MI and major bleeding was defined per trial definition. ST was defined as per Academic Research Consortium (ARC) [22].

The statistical analysis was performed according to the recommendations from the Cochrane Collaboration using Review Manager Version 5.1 (The Nordic Cochrane Center, The Cochrane Collaboration, 2008, Copenhagen). A random-effects model with inverse variance weighting was used to calculate odds ratio (OR) and 95% confidence interval, associated with S-DAPT versus L-DAPT for above end points. Forest plots were used to observe the overall effect of studies for each endpoint. Heterogeneity between studies was assessed using Cochrane's Q test and I^2 statistic, which denotes the percentage of total variation across studies that is a result of heterogeneity rather than chance. $I^2 < 25\%$ was considered as low heterogeneity and $I^2 > 75\%$ as high. Heterogeneity was considered significant if the p value was less than 0.05. Publication bias was assessed by Begg's test and Egger's regression test.

RESULTS:

2819 publications were found at initial search and after abstract and manuscript evaluation eight studies are selected for final analysis (Figure 1). Investigators of OPTIMIZE and OPTIDUAL trials have shared their data on outcomes for DM and non-DM patients after PCI for this analysis, which was not reported/published before [5, 8]. In the selected eight RCTs, 28,318 patients were randomized to S-DAPT (N=14,095) versus L-DAPT (14, 203) (Table 1) [4-11]. Out of 28,318 patients 8,234 had DM and 20,084 were non-diabetics. Median follow up period in studies included in our analysis varied from 9 months to 33 months with overall median follow up of 12 months after

completion of L-DAPT. Use of second generation DES varied from 36% to 100%.

Baseline characteristics of patients and studies included in the final analysis are summarized in Table 2.

On analysis, of 18025 patients (5417 DM and 12608 non-DM), S-DAPT was associated with an increased rate of ST in non-DM patients [3.67 (2.04, 6.59), $p<0.0001$] (Figure2). There was no significant difference in the rate of ST with S-DAPT among DM patients [1.52 (0.70, 3.29), $p=0.29$] (Figure2) (Table 3). There was borderline interaction between subgroups for ST ($P_{\text{interaction}}=0.07$). Overall, regardless of DAPT duration, DM patients have higher rates of ST (0.72%) than non-DM patients (0.57%). There was also no significant difference in the rate of MI with S-DAPT and L-DAPT in DM [1.33 (0.88, 2.01), $p=0.18$] and non-DM [1.52 (0.81, 2.83), $p=0.19$] patients on analysis of 18025 patients (5417 DM and 12608 non-DM) (Figure 3) (Table 3). DM patients have higher rates of MI (3.30%) than non-DM (2.08%) irrespective of the duration of DAPT.

There was also no significant difference in the rate of all cause mortality, and cardiac mortality with S-DAPT and L-DAPT in DM [1.19 (0.72-1.95) and 1.25 (0.69, 2.25) respectively] and non-DM-patients [0.93 (0.58, 1.48); and 0.75 (0.42, 1.35) respectively] (Figure 4, 5) (Table 3). However, overall event rates of both all-cause mortality (3.16% vs 1.89%), and cardiac mortality (2.01% vs 0.83%) were higher for DM patients as compared to non-DM patients.

Similarly, no significant difference was found between S-DAPT and L-DAPT for TVR, major bleeding and stroke for DM [1.39 (0.89, 2.17); 0.92 (0.19, 4.42); and 0.98 (0.29, 3.28) respectively] and non-DM patients [0.99 (0.71, 1.39); 0.72 (0.28, 1.84); and 1.01 (0.40, 2.56) respectively] (Supplementary Figure 1, 2, 3) (Table 3). On analysis of 16,980 patients (4762 with DM and 12218 non-DM patients) there was no significant difference in the rate of composite end point-NACE with S-DAPT and L-DAPT for DM [0.94 (0.57, 1.54)] and non-DM patients [1.01 (0.77, 1.32)] (Supplementary figure 4). However, overall event rates were higher among DM patients compared to non-DM (4.49% vs 3.01%).

DISCUSSION:

In our meta-analysis of RCTs in which patients were analyzed according to DM status, S-DAPT compared to L-DAPT was associated with a significant increase in rate of ST in non-DM patients. However, there was no significant difference in the rate of all cause mortality, cardiac mortality, MI, ST and TVR with S-DAPT among DM patients.

Although widespread use of DES has improved overall clinical outcomes [23, 24], sustained risk of ST and MI remain a major concern in DM patients [25-27]. This elevated risk has been partly attributed to existence of a pro-thrombotic state in DM patients secondary to several factors, including endothelial dysfunction, hypercoagulation and platelet hyperactivation [28, 29]. Furthermore, patients with impaired glucose metabolism have been reported to have a suboptimal biochemical response to

aspirin (a mainstay agent for DAPT)—‘aspirin resistance’ [30, 31]. Possible causes of this suboptimal response have been proposed including aspirin formulation used and dose of aspirin given [32]. Recently, in a randomized, single-blind, triple-crossover study Bhatt et al, investigated if suboptimal response to aspirin in diabetic patients is mediated via decrease oral bioavailability [33]. Forty patients with type 2 DM and without cardiovascular disease were randomized to receive a plain aspirin tablet, a PL2200 aspirin capsule (modified-release lipid-based aspirin), and a delayed-release enteric-coated (EC) aspirin caplet. There was no significant difference in aspirin response between plain aspirin and PL2200 aspirin ($p=0.41$), whereas aspirin response was quicker with the former two when compared to EC aspirin ($p<0.001$). The patients on EC aspirin were found to have lower levels of plasma aspirin, secondary to decreased absorption. Therefore, use of non-EC aspirin formulation might be more appropriate in this patient population.

DAPT has been shown to be protective against early and late ischemic events after DES implantation, although the optimal duration remains unclear [12-20]. Even though DM patients have higher prevalence of CAD and they undergo coronary revascularization procedures, including PCI, more often than non-DM patients, there is paucity of data evaluating the impact of DM on duration of DAPT [34]. DM has been considered an independent variable in DAPT score, presence of which might favor L-DAPT after PCI [35]. However, in recent patient-level pooled analysis of 11,473 patients, Gargiulo and colleagues have reported that L-DAPT was associated with increased bleeding

complications in both DM and non-DM patients, without significant reduction in ischemic events [36].

In this context, our meta-analysis of RCTs provides important insights into relationship of DAPT duration with clinical outcomes among DM and non-DM patients after implantation of DES. Our results suggest that DM patients have higher rate of adverse CV events than non-DM patients after PCI with DES implantation irrespective of DAPT duration. Further, L-DAPT compared to S-DAPT might not be effective in reducing adverse cardiovascular outcomes after PCI in DM patients.

While placing our results in the context of above findings, several points are worth mentioning. First, accelerated neointimal hyperplasia leading to endothelialization may account for the differential effect of prolonged DAPT on ST in patients with DM [37, 38]. Second, diminished impact of antiplatelet agents in DM, especially clopidogrel, on platelet inhibition may explain a higher incidence of ST in the first 6 months, as seen in S-DAPT arm of EXCELLENT trial, where the reported ST events occurred while the patients were still on DAPT [7, 39]. Third, platelet hyper-reactivity has been implicated as a cause of increased thrombogenesis in DM, suggesting a place for more potent anti-platelets [40, 41]. Although, relatively low number of events and inclusion of studies with short-term follow up [7, 10] precludes definitive conclusions, our results suggest complex interaction of various clinical end points and DM after DES implantation.

DM has been shown to be an independent risk factor for ST [42]. One of the major issues surrounding DAPT duration is a sustained risk of late (1-12 months) [43] and very late (>12 months) ST [44] after DES implantation. Several medium-sized RCTs have shown that S- DAPT is non-inferior to L- DAPT in terms of incidence of ST [5, 6, 10], however these studies did not possess sufficient statistical power to assess ST. The Dual Antiplatelet Therapy (DAPT) study, the only study adequately powered for assessing DAPT duration on rates of ST, randomized event-free patients at 12 months to placebo versus extended (30 months) duration of DAPT and found a significantly reduced risk of ST with the L-DAPT (0.4% vs 1.4%, $p < 0.001$) [22]. However, consistent with our finding, the protective effect of L-DAPT in DAPT trial was attenuated in DM patients (0.6% vs 1.1%), compared to non -DM (0.3% vs 1.5%) [11]. Similarly, in analysis of the diabetic subgroup of the EXCELLENT (Efficacy of Xience/Promus Versus Cypher to Reduce Late Loss After Stenting) trial, there was trend toward increased risk of ST with S-DAPT when compared with standard 12-month DAPT [7]. Our analysis demonstrated significantly increased risk of ST with S-DAPT relative to L-DAPT in non-diabetic patients. Although in DM patients there was no statistically significant difference in the rate of ST with L-DAPT as compared to S-DAPT. This could be partly due to relatively small number of patients and events in the DM arm (5417 patients and 39 ST) compared to non-DM (12608 patients and 72 ST). This could be due to higher baseline platelet activity in DM patients, which remains higher than non-DM even after DAPT.

Our analysis has several limitations. First, this is a meta-analysis performed on study-level data due to non-availability of patient-level data. Thus, covariate-adjusted analyses

for possible confounders like generation of DES, clinical presentation/indication of PCI, location and complexity of lesions during PCI, baseline clinical characteristics (e.g. insulin dependent or non-insulin dependent DM, renal dysfunction, heart failure) could not be performed. Similarly, due to lack of patient-level information regarding use of statins, beta-blockers or renin–angiotensin system blockers, we could not evaluate their impact on the relative efficacy of prolonged DAPT in DM and non-DM patients. Studies included in our meta-analysis randomized patients with various clinical presentations and characteristics, thus it is unlikely that these variables would have confounded our results due to selection bias; however, this cannot be completely excluded on the basis of current analysis. Second, prasugrel and ticagrelor have been shown to be associated with decrease in the rate of ST compared to clopidogrel and have recently been recommended by the American College of Cardiology/American Heart Association and European Society of Cardiology for management of ACS [45, 46]. In the present study, we were unable to evaluate duration of DAPT with prasugrel and ticagrelor, as studies included in our analysis did not compare duration of DAPT using these antiplatelets separately. However, such variations in DAPT reflect real world clinical practice, where different second antiplatelet agents are selected based on operator choices, clinical settings and drug availability. Third, trials included in our analysis were open label and timing of randomization was not the same in all studies, which could dilute the difference between the S-DAPT and L-DAPT arms, especially if ischemic or bleeding events have occurred before randomization or DAPT discontinuation.

CONCLUSIONS:

Compared to L-DAPT, S-DAPT was associated with significant increase in rate of ST in non-DM patients. Duration of DAPT had no significant impact on rates of all cause mortality, cardiac mortality, MI, ST and TVR among DM patients.

REFERENCES:

1. Hammoud T, Tanguay JF, Bourassa MG. Management of coronary artery disease: therapeutic options in patients with diabetes. *J Am Coll Cardiol.* 2000 ;36:355-65.
2. The BARI investigators. Influence of diabetes on 5-year mortality and morbidity in a randomized trial comparing CABG and PTCA in patients with multivessel disease: the Bypass Angioplasty Revascularization Investigation (BARI). *Circulation.* 1997; 96:1761-9.
3. Montalescot G, Brieger D, Dalby AJ, Park SJ, Mehran R. Duration of Dual Antiplatelet Therapy After Coronary Stenting: A Review of the Evidence. *J Am Coll Cardiol.* 2015; 66:832-47.
4. Kim BK, Hong MK, Shin DH, Nam CM, Kim JS, Ko YG, Choi D, Kang TS, Park BE, Kang WC, Lee SH, Yoon JH, Hong BK, Kwon HM, Jang Y; RESET Investigators. A new strategy for discontinuation of dual antiplatelet therapy: the RESET Trial (REal Safety and Efficacy of 3-month dual antiplatelet Therapy following Endeavor zotarolimus-eluting stent implantation). *J Am Coll Cardiol.* 2012 ;60:1340-8.
5. Feres F, Costa RA, Abizaid A, Leon MB, Marin-Neto JA, Botelho RV, King SB 3rd, Negoita M, Liu M, de Paula JE, Mangione JA, Meireles GX, Castello HJ Jr, Nicolela EL Jr, Perin MA, Devito FS, Labrunie A, Salvadori D Jr, Gusmão M, Staico R, Costa JR Jr,

de Castro JP, Abizaid AS, Bhatt DL; OPTIMIZE Trial Investigators. Three vs twelve months of dual antiplatelet therapy after zotarolimus-eluting stents: the OPTIMIZE randomized trial. *JAMA*. 2013; 310:2510-22.

6. Schulz-Schüpke S, Byrne RA, Ten Berg JM, Neumann FJ, Han Y, Adriaenssens T, Tölg R, Seyfarth M, Maeng M, Zrenner B, Jacobshagen C, Mudra H, von Hodenberg E, Wöhrle J, Angiolillo DJ, von Merzljak B, Rifatov N, Kufner S, Morath T, Feuchtenberger A, Ibrahim T, Janssen PW, Valina C, Li Y, Desmet W, Abdel-Wahab M, Tiroch K, Hengstenberg C, Bernlochner I, Fischer M, Schunkert H, Laugwitz KL, Schömig A, Mehilli J, Kastrati A; Intracoronary Stenting and Antithrombotic Regimen: Safety And Efficacy of 6 Months Dual Antiplatelet Therapy After Drug-Eluting Stenting (ISAR-SAFE) Trial Investigators. ISAR-SAFE: a randomized, double-blind, placebo controlled trial of 6 vs. 12 months of clopidogrel therapy after drug-eluting stenting. *Eur Heart J*. 2015 ;36:1252-63.

7. Gwon HC, Hahn JY, Park KW, Song YB, Chae IH, Lim DS, Han KR, Choi JH, Choi SH, Kang HJ, Koo BK, Ahn T, Yoon JH, Jeong MH, Hong TJ, Chung WY, Choi YJ, Hur SH, Kwon HM, Jeon DW, Kim BO, Park SH, Lee NH, Jeon HK, Jang Y, Kim HS. Six month versus 12-month dual antiplatelet therapy after implantation of drug-eluting stents: the Efficacy of Xience/Promus Versus Cypher to Reduce Late Loss After Stenting (EXCELLENT) randomized, multicenter study. *Circulation*. 2012 ;125:505-13.

8. Helft G, Steg PG, Le Feuvre C, Georges JL, Carrie D, Dreyfus X, Furber A, Leclercq

F, Eltchaninoff H, Falquier JF, Henry P, Cattan S, Sebagh L, Michel PL, Tuambilangana A, Hammoudi N, Boccara F, Cayla G, Douard H, Diallo A, Berman E, Komajda M, Metzger JP, Vicaute E; OPTImal DUAL Antiplatelet Therapy Trial Investigators.

Stopping or continuing clopidogrel 12 months after drug-eluting stent placement: the OPTIDUAL randomized trial. *Eur Heart J.* 2016 ;37:365-74

9. Collet JP, Silvain J, Barthélémy O, Rangé G, Cayla G, Van Belle E, Cuisset T, Elhadad S, Schiele F, Lhoest N, Ohlmann P, Carrié D, Rousseau H, Aubry P, Monségu J, Sabouret P, O'Connor SA, Abtan J, Kerneis M, Saint-Etienne C, Beygui F, Vicaute E, Montalescot G; ARCTIC investigators. Dual-antiplatelet treatment beyond 1 year after drug-eluting stent implantation (ARCTIC-Interruption): a randomised trial. *Lancet.* 2014 ; 384:1577-85.

10. Lee CW, Ahn JM, Park DW, Kang SJ, Lee SW, Kim YH, Park SW, Han S, Lee SG, Seong IW, Rha SW, Jeong MH, Lim DS, Yoon JH, Hur SH, Choi YS, Yang JY, Lee NH, Kim HS, Lee BK, Kim KS, Lee SU, Chae JK, Cheong SS, Suh IW, Park HS, Nah DY, Jeon DS, Seung KB, Lee K, Jang JS, Park SJ. Optimal duration of dual antiplatelet therapy after drug-eluting stent implantation: a randomized, controlled trial. *Circulation.* 2014;129:304-312.

11. Mauri L, Kereiakes DJ, Yeh RW, Driscoll-Shempp P, Cutlip DE, Steg PG, Normand SL, Braunwald E, Wiviott SD, Cohen DJ, Holmes DR Jr, Krucoff MW, Hermiller J, Dauerman HL, Simon DI, Kandzari DE, Garratt KN, Lee DP, Pow TK, Ver Lee P,

Rinaldi MJ, Massaro JM; DAPT Study Investigators. Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *N Engl J Med*. 2014; 371:2155-66.

12. Giustino G, Baber U, Sartori S, Mehran R, Mastoris I, Kini AS, Sharma SK, Pocock SJ, Dangas GD. Duration of dual antiplatelet therapy after drug-eluting stent implantation: a systematic review and meta-analysis of randomized controlled trials. *J Am Coll Cardiol*. 2015;65:1298-310

13. Navarese EP, Andreotti F, Schulze V, Kołodziejczak M, Buffon A, Brouwer M, Costa F, Kowalewski M, Parati G, Lip GY, Kelm M, Valgimigli M. Optimal duration of dual antiplatelet therapy after percutaneous coronary intervention with drug eluting stents: meta-analysis of randomised controlled trials. *BMJ*. 2015; 350:h1618

14. Palmerini T, Benedetto U, Bacchi-Reggiani L, Della Riva D, Biondi-Zoccai G, Feres F, Abizaid A, Hong MK, Kim BK, Jang Y, Kim HS, Park KW, Genereux P, Bhatt DL, Orlandi C, De Servi S, Petrou M, Rapezzi C, Stone GW. Mortality in patients treated with extended duration dual antiplatelet therapy after drug-eluting stent implantation: a pairwise and Bayesian network meta-analysis of randomised trials. *Lancet*. 2015; 385:2371-82.

15. Elmariah S, Mauri L, Doros G, Galper BZ, O'Neill KE, Steg PG, Kereiakes DJ, Yeh RW. Extended duration dual antiplatelet therapy and mortality: a systematic review and meta-analysis. *Lancet*. 2015;385:792-8.

16. El-Hayek G, Messerli F, Bangalore S, Hong MK, Herzog E, Benjo A, Tamis-Holland JE. Meta-analysis of randomized clinical trials comparing short-term versus long-term dual antiplatelet therapy following drug-eluting stents. *Am J Cardiol.* 2014; 114:236-42.
17. Sharma A, Sharma SK, Vallakati A, Garg A, Lavie CJ, Mukherjee D, Marmur JD. Duration of dual antiplatelet therapy after various drug-eluting stent implantation. *Int J Cardiol.* 2016 Jul 15;215:157-66.
18. Sharma A, Lavie CJ, Sharma SK, Garg A, Vallakati A, Mukherjee D, Marmur JD. Duration of Dual Antiplatelet Therapy After Drug-Eluting Stent Implantation in Patients With and Without Acute Coronary Syndrome: A Systematic Review of Randomized Controlled Trials. *Mayo Clin Proc.* 2016 Aug;91(8):1084-93.
19. Sharma A, Hai O, Garg A, Vallakati A, Lavie CJ, Marmur JD. Duration of Dual Antiplatelet Therapy Following Drug-Eluting Stent Implantation: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Current Problems in Cardiology: In press.*
20. Palmerini T, Sangiorgi D, Valgimigli M, Biondi-Zoccai G, Feres F, Abizaid A, Costa RA, Hong MK, Kim BK, Jang Y, Kim HS, Park KW, Mariani A, Della Riva D,

Genereux P, Leon MB, Bhatt DL, Bendetto U, Rapezzi C, Stone GW. Short- versus longterm dual antiplatelet therapy after drug-eluting stent implantation: an individual patient data pairwise and network meta-analysis. *J Am Coll Cardiol*. 2015 ;65:1092-102.

21. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009 ;151:264-9, W64.

22. Cutlip DE, Windecker S, Mehran R, et al. Clinical End Points in Coronary Stent Trials: A Case for Standardized Definitions. *Circulation*. 2007 May 1;115(17):2344–51.

23. Palmerini T, Benedetto U, Biondi-Zoccai G, Della Riva D, et al. Long-Term Safety of Drug-Eluting and Bare-Metal Stents: Evidence From a Comprehensive Network MetaAnalysis. *J Am Coll Cardiol*. 2015 ;65:2496-507.

24. Bangalore S, Kumar S, Fusaro M, Amoroso N, et al. Short- and long-term outcomes with drug-eluting and bare-metal coronary stents: a mixed-treatment comparison analysis of 117 762 patient-years of follow-up from randomized trials. *Circulation*. 2012;125:2873-91.

25. Simsek C, Räber L, Magro M, Boersma E, et al. Long-term outcome of the unrestricted use of everolimus-eluting stents compared to sirolimus-eluting stents and

paclitaxel-eluting stents in diabetic patients: the Bern-Rotterdam diabetes cohort study.

Int J Cardiol. 2013 ;170:36-42.

26. Cassese S, Byrne RA, Tada T, Pinieck S, et al. Incidence and predictors of restenosis after coronary stenting in 10 004 patients with surveillance angiography. Heart. 2014; 100:153-9.

27. Moussa I, Leon MB, Baim DS, O'Neill WW, et al. Impact of sirolimus-eluting stents on outcome in diabetic patients: a SIRIUS (SIRolImUS-coated Bx Velocity balloonexpandable stent in the treatment of patients with de novo coronary artery lesions) substudy. Circulation. 2004 ;109:2273-8.

28. Osende JJ, Badimon JJ, Fuster V, Herson P, Rabito P, Vidhun R, Zaman A, Rodriguez OJ, Lev EI, Rauch U, Heflt G, Fallon JT, Crandall JP. Blood thrombogenicity in Type 2 diabetes mellitus patients is associated with glycemic control. J Am Coll Cardiol. 2001 ;38:1307-12.

29. Neergaard-Petersen S, Hvas AM, Kristensen SD, Grove EL, Larsen SB, Phoenix F, Kurdee Z, Grant PJ, Ajjan RA. The influence of type 2 diabetes on fibrin clot properties in patients with coronary artery disease. Thromb Haemost. 2014; 112:1142-50.

30. Smith JP, Haddad EV, Taylor MB, et al. Suboptimal inhibition of platelet cyclooxygenase-1 by aspirin in metabolic syndrome. Hypertension 2012;59:719-25.

31. DiChiara J, Bliden KP, Tantry US, et al. The effect of aspirin dosing on platelet function in diabetic and nondiabetic patients: an analysis from the aspirin-induced platelet effect (ASPECT) study. *Diabetes* 2007;56:3014-9.
32. Ertugrul DT, Tural E, Yildiz M, et al. Aspirin resistance is associated with glycemic control, the dose of aspirin, and obesity in type 2 diabetes mellitus. *J Clin Endocrinol Metab* 2010;95:2897-901.
33. Bhatt DL, Grosser T, Dong JF, et al. Enteric Coating and Aspirin Nonresponsiveness in Patients With Type 2 Diabetes Mellitus. *J Am Coll Cardiol* 2017;69:603-12.
34. Flaherty JD, Davidson CJ. Diabetes and coronary revascularization. *JAMA*. 2005; 293:1501-8.
35. Yeh RW, Secemsky EA, Kereiakes DJ, Normand SL, Gershlick AH, Cohen DJ, Spertus JA, Steg PG, Cutlip DE, Rinaldi MJ, Camenzind E, Wijns W, Apruzzese PK, Song Y, Massaro JM, Mauri L; DAPT Study Investigators. Development and Validation of a Prediction Rule for Benefit and Harm of Dual Antiplatelet Therapy Beyond 1 Year After Percutaneous Coronary Intervention. *JAMA*. 2016 ;315:1735-49.
36. Gargiulo G, Windecker S, da Costa BR, Feres F, Hong MK, Gilard M, Kim HS, Colombo A, Bhatt DL, Kim BK, Morice MC, Park KW, Chieffo A, Palmerini T, Stone

GW, Valgimigli M. Short term versus long term dual antiplatelet therapy after implantation of drug eluting stent in patients with or without diabetes: systematic review and meta-analysis of individual participant data from randomised trials. *BMJ*. 2016; 355:i5483.

37. Luscher TF, Creager MA, Beckman JA, Cosentino F. Diabetes and vascular disease: Pathophysiology, clinical consequences, and medical therapy: Part II. *Circulation*. 2003; 108:1655–1661.

38. Brar SS, Kim J, Brar SK, Zadegan R, et al. Long-Term Outcomes by Clopidogrel Duration and Stent Type in a Diabetic Population With De Novo Coronary Artery Lesions. *J Am Coll Cardiol* 2008;51:2220–7.

39. Serebruany V, Pokov I, Kulickowski W, Chesebro J, Badimon J. Baseline platelet activity and response after clopidogrel in 257 diabetics among 822 patients with coronary artery disease. *Thromb Haemost* 2008;100:76–82.

40. Ferreira JL, Angiolillo DJ. Diabetes and antiplatelet therapy in acute coronary syndrome. *Circulation* 2011;123:798-813.

41. Wiviott SD, Braunwald E, Angiolillo DJ, Meisel S, et al. Greater clinical benefit of more intensive oral antiplatelet therapy with prasugrel in patients with diabetes mellitus in the trial to assess improvement in therapeutic outcomes by optimizing platelet

inhibition with prasugrel- thrombolysis in myocardial infarction 38. *Circulation*. 2008; 118:1626–1636.

42. Iakovou I, Schmidt T, Bonizzoni E, Ge L, et al. Incidence, predictors, and outcome of thrombosis after successful implantation of drug-eluting stents. *JAMA*. 2005;293:2126 –2130.

43. Yano M, Natsuaki M, Morimoto T, Nakagawa Y, et al. Antiplatelet therapy discontinuation and stent thrombosis after sirolimus-eluting stent implantation: Five-year outcome of the j-Cypher Registry. *Int J Cardiol*. 2015 Jun 18;199:296-301.

44. Armstrong EJ, Maddox TM, Carey EP, Grunwald GK, et al. Mortality after presentation with stent thrombosis is associated with time from index percutaneous coronary intervention: a report from the VA CART program. *Am Heart J*. 2014; 168:560-7.

45. Amsterdam EA, Wenger NK, Brindis RG, Casey DE Jr, Ganiats TG, Holmes DR Jr, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ; American College of Cardiology; American Heart Association Task Force on Practice Guidelines; Society for Cardiovascular Angiography and Interventions; Society of Thoracic Surgeons; American Association for Clinical Chemistry. 2014 AHA/ACC Guideline for the Management of Patients with Non-ST-Elevation Acute Coronary Syndromes: a report of the American

College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. 2014 ;64:e139-228.

46. Authors/Task Force Members, Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, Bax JJ, Borger MA, Brotons C, Chew DP, Gencer B, Hasenfuss G, Kjeldsen K, Lancellotti P, Landmesser U, Mehilli J, Mukherjee D, Storey RF, Windecker S. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). Eur Heart J. 2016 ;37:267-315.

FIGURES LEGENDS:

Figure 1 The Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) flow chart for the trial selection process

Figure 2: Forest plot for stent thrombosis for DM and non-DM patients

Figure 3: Forest plot for myocardial infarction for DM and non-DM patients

Figure 4. Forest plot for all cause mortality for DM and non-DM patients

Figure 5. Forest plot for cardiac mortality for DM and non-DM patients

Tables:

Table 1. Baseline characteristics of studies included in analysis

Table 2. Baseline patient characteristics in the studies included in the analysis

Table 3. Outcomes with S-DAPT and L-DAPT in DM and non-DM patients

STUDY	DESIGN	Time to Randomization	Inclusion Criteria	Exclusion Criteria	Primary end-point
RESET (2012)	Randomized, open label, multicentre	At index PCI	20-85 years old; $\geq 50\%$ stenosis; elective PCI; stable angina, unstable angina, or acute MI	CI to anti-platelets; STEMI within 48 h; CVA; PAD; thromboembolism; ST; ISR; CTO; LM stenosis $> 50\%$; History of DES Implantation; cardiogenic shock.	Composite of Cardiac Death, MI, ST, ischemia-driven TVR or Bleeding at 12 Months after PCI
OPTIMIZE (2014)	Randomized, open label, multicentre	At index PCI	Stable angina or silent ischemia or low risk ACS as defined by unstable angina or recent (but not acute) myocardial infarction (< 30 days)	STEMI for Primary or Rescue PCI; BMS less than 6 months prior to index procedure; previous treatment with DES; scheduled surgery within 12 months; CI to anti-platelets; ISRs	Composite of Death, MI, Stroke or Major Bleeding at 12 Months after PCI
ISAR-SAFE (2014)	Randomized, double-blind,	6 months after Index PCI	Patients on clopidogrel at 6	Previous ST; DES in LM; MI in the previous 6 months; Planned major	Composite of Death, MI, Stroke or ST or TIMI major

	multicentr e		(-1/+2) months after PCI with DES	surgery within next 6 months; CI to anti- platelets; OAC; Prior ICH	Bleeding at 15 Months after PCI
EXCELLENT (2012)	Randomiz ed, open label, multicentr e	At index PCI	≥1 lesion native coronary vessel; >50% stenosis; stable angina, unstable angina, recent MI, silent ischemia, positive function al study, or reversibl e changes on ECG consiste nt with ischemia	MI within 72 hours; any prior stent implanted in the target vessel; cardiogenic shock; major bleeding within 3 months; elective surgical procedure within 12 months; left main stenosis > 50%; CTOs; bifurcation lesions requiring 2- stent strategy.	Composite of Cardiac Death, MI or TVR at 12 Months after PCI
OPTIDUAL (2015)	Randomiz ed, open label, multicentr e	12 months after index PCI	Stable angina, silent ischaemi a, or acute coronary syndrom e with ≥1 lesion	Requirement for oral anticoagulation , DES implantation in an unprotected left main coronary artery, and	Composite of all-cause mortality, non-fatal myocardial infarction, stroke, or major bleeding.

			with stenosis >50% located in a native vessel implanted with ≥1 DES of any type.	malignancies or other coexisting conditions associated with a life expectancy of <2 years	
ARCTIC (2014)	Randomized, open label, multicentre	12 months after index PCI	≥18 years and eligible for PCI with planned use of ≥1 DES.	Primary PCI for STEMI; chronic anticoagulation or bleeding diathesis; CI to APT; active bleeding or major surgery within 3 months; scheduled surgery within 1 year	Composite of Death, MI, Stroke or TIA, urgent revascularization or ST
DES-LATE (2014)	Randomized, open label, multicentre	12 months after index PCI	DES within 12 months; on DAPT; no MACE (MI, stroke, repeat PCI) or major bleeding since PCI.	CI to antiplatelet drugs, concomitant vascular disease or recent ACS requiring clopidogrel use	Composite of Cardiac Death, MI or Stroke at 24 Months after PCI
DAPT	Randomiz	12 months	>18	Planned major surgery within	ST; Composite of

(2014)	ed, double- blind, multicentr e	after index PCI	years undergoi ng FDA- approve d PCI with DES or BMS; No MACE or bleeding within 12 months after procedur e.	the 30 months post enrollment; OAC; Both BMS and DES during index procedure; PCI or surgery between 6 weeks post-PCI and randomization	death, MI, stroke; and Moderate or severe GUSTO bleeding
--------	---	--------------------	---	---	---

Table 1. Baseline characteristics of studies included in analysis

ACS: Acute coronary syndrome, BMS: bare metal stent, CI: contraindication; CKD: Chronic kidney disease, CTO: Chronic total occlusion, ECG: Electrocardiogram, FDA: Food and Drug Administration, LM: Left main, TVR: target vessel revascularization, STEMI: ST segment elevation myocardial infarction, NSTEMI: non ST segment elevation myocardial infarction, PCI: Percutaneous coronary interventions, MI: myocardial infarction, DES, drug eluting stents, OAC: oral anticoagulant, MACE: major adverse cardiac event, GUSTO: Global Utilization Of Streptokinase And Tpa For Occluded Arteries, SVG: Saphenous venous grafts, UPLMA: Unprotected left main artery.

ARCTIC: Assessment by a double Randomisation of a Conventional antiplatelet strategy versus a monitoring-guided strategy for drug-eluting stent implantation and, of Treatment Interruption versus Continuation 1 year after stenting; DAPT: The dual antiplatelet therapy study; DES-LATE: the Optimal Duration of Clopidogrel Therapy With DES to Reduce Late Coronary Arterial Thrombotic Event; EXCELLENT: Efficacy of Xience/Promus Versus Cypher in rEducing Late Loss After stENTing; ISAR-SAFE: Safety and Efficacy of Six Months Dual Antiplatelet Therapy After Drug-Eluting Stenting; OPTIDUAL: Optimal Duration of Dual Antiplatelet Therapy After Drug-eluting Stent Implantation; OPTIMIZE: Optimized Duration of Clopidogrel Therapy Following Treatment With the Zotarolimus-Eluting Stent in Real-World Clinical Practice; RESET: REal Safety and Efficacy of 3-month dual antiplatelet Therapy following Endeavor zotarolimus-eluting stent implantation).

Study	N Total	N DM	N Non- DM	Follow -up (months)	Age (y)	Male (%)	HT N (%)	HL D (%)	Prior MI (%)	ACS (%)	2 nd gen ste nt (%)
RESET	2117	3 months s=146	3 months = 913	12	62	64	62	59	2	28	85
		12 months s=146	12 months = 912								
OPTIMIZE	3119	3 months s=554	3 months = 1009	12	62	63	87	64	35	32	100
		12 months s=549	12 months = 1007								
ISAR-SAFE	3997	6 months s=495	6 months = 1501	9	67	81	90	87	25	40	89
		12 months s=484	12 months = 1517								
EXCELLENT	1443	6 months s=272	6 months = 450	12	63	65	73	76	5	51	75

		12 month s= 278	12month hs= 443								
ARCTIC	125 9	12 month s= 222	12 months = 402	17	64	80	60	67	30	26	63
		18 month s= 198	18 months = 437								
DES- LATE	504 5	12 month s= 709	12 months = 1805	24	62	69	57	NA	4	61	36
		30 month s= 709	30 months = 1822								
DAPT	996 1	12 month s= 1481	12 months = 3460	18	62	75	75	NA	22	26	61
		30 month s= 1556	30 months = 3464								
OPTIDU AL	138 5	12 month s= 222	12 months = 468	33	64	81	59	NA	17	36	66
			48								

48	months
month	=
s=	482
213	

Table 2. Baseline patient characteristics in the studies included in the analysis

?: percent; ACS: Acute coronary syndrome, N: Number of patients, DM: diabetes mellitus, MI: myocardial infarction, HTN: hypertension, HLD: hyperlipidemia

ARCTIC: Assessment by a double Randomisation of a Conventional antiplatelet strategy versus a monitoring-guided strategy for drug-eluting stent implantation and, of Treatment Interruption versus Continuation 1 year after stenting; DAPT: The dual antiplatelet therapy study; DES-LATE: the Optimal Duration of Clopidogrel Therapy With DES to Reduce Late Coronary Arterial Thrombotic Event; EXCELLENT: Efficacy of Xience/Promus Versus Cypher in rEducing Late Loss After stENTing; ISAR-SAFE: Safety and Efficacy of Six Months Dual Antiplatelet Therapy After Drug-Eluting Stenting; OPTIDUAL: Optimal Duration of Dual Antiplatelet Therapy After Drug-eluting Stent Implantation; OPTIMIZE: Optimized Duration of Clopidogrel Therapy Following Treatment With the Zotarolimus-Eluting Stent in Real-World Clinical Practice; RESET: REal Safety and Efficacy of 3-month dual antiplatelet Therapy following Endeavor zotarolimus-eluting stent implantation).

	All patients (N=28,318)	Diabetics (N=8,234)	Non- Diabetics (N=20,084)	P_{interaction}	Hetero- geneity (I²)
All cause mortality	1.04 [0.74, 1.46]	1.19 [0.72-1.95]	0.93 [0.58, 1.48]	0.48	0%
Cardiac mortality	0.97 [0.64, 1.46]	1.25 [0.69, 2.25]	0.75 [0.42, 1.35]	0.23	0%
MI	1.48 [1.00, 2.19]	1.33 [0.88, 2.01]	1.52 [0.81, 2.83]	0.73	58%
ST	2.16 [1.21, 3.85]	1.52 [0.70, 3.29]	3.67 [2.04, 6.59]	0.07	21%
TVR	1.12 [0.85, 1.47]	1.39 [0.89, 2.17]	0.99 [0.71, 1.39]	0.24	20%
Major bleeding*	0.77 [0.34, 1.72]	0.92 [0.19, 4.42]	0.72 [0.28, 1.84]	0.79	0%
Stroke	1.00 [0.48, 2.09]	0.98 [0.29, 3.28]	1.01 [0.40, 2.56]	0.98	0%
NACE	0.97 [0.76, 1.25]	0.94 [0.57, 1.54]	1.01 [0.77, 1.32]	0.81	48%

S-DAPT compared with L-DAPT in diabetics and non-diabetics; *: GUSTO or TIMI major bleeding

MI: myocardial infarction; NACE: net adverse clinical events; ST: Stent thrombosis (definite or probable); TVR: target vessel revascularization

Conflict of interest: None.

ACCEPTED MANUSCRIPT




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9